


That which is claimed is:

1. A method for modulating process(es) mediated by peroxisome proliferator activated receptor-gamma (PPAR- γ), said method comprising conducting said process(es) in the presence of at least one PPAR- γ -selective prostaglandin or prostaglandin-like compound or precursor thereof.

2. A method according to Claim 1 wherein said PPAR- γ -selective prostaglandin is selected from a prostaglandin- J_2 , a prostaglandin- D_2 , or a precursor thereof.

3. A method according to Claim 2 wherein said prostaglandin- J_2 is selected from prostaglandin- J_2 , Δ^{12} -prostaglandin- J_2 or 15-deoxy- $\Delta^{12,14}$ -prostaglandin- J_2 .

4. A method according to Claim 1, wherein said PPAR- γ -selective prostaglandin or prostaglandin-like compound has the structure I:



(I)

wherein:

A is selected from hydrogen or a leaving group at the α - or β - position of the ring, or A is absent when there is a double bond between C^α and C^β of the ring;

X is an alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or

20 substituted alkynyl group having in the range of 2 up to 15 carbon atoms; and Y is an alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl group having in the range of 2 up to 15 carbon atoms.

5. A method according to claim 4 wherein:





X of Formula I is selected from:

- 5
- (CRR)_m-Z,
 - (CRR)_m-C(R)=C(R)-(CRR)_{m'}-Z, or
 - 5 -(CRR)_m-C≡C-(CRR)_{m''}-Z, wherein:
- each R is independently selected from hydrogen, lower alkyl, substituted lower alkyl, hydroxy, lower alkoxy, thioalkyl, halogen, trifluoromethyl, 10 cyano, nitro, amino, carboxyl, carbamate, sulfonyl or sulfonamide, m falls in the range of 1 up to 15, each m' falls independently in the range of 0 up to 12, with the proviso that 15 the total chain length of the alkenyl moiety does not exceed 15 carbon atoms, each m'' falls independently in the range of 0 up to 12, with the proviso that 20 the total chain length of the alkynyl moiety does not exceed 15 carbon atoms, and Z is a polar, heteroatom-containing substituent; and


09455552.092498

25

Y of Formula I is selected from:



30



wherein

each R is independently as defined
above,

35

each R' is independently selected
from H, lower alkyl, substituted
lower alkyl, or a leaving group,

Z' is selected from H, lower alkyl or
substituted lower alkyl,

n falls in the range of 0 up to 4,

40

n' falls in the range of 2 up to 12, and

n'' falls in the range of 1 up to 3.

6. A method according to claim 5 wherein Z is
selected from cyano, nitro, amino, carbamate, or a
substituent having the structure:

5

$-CH_2OR'$, wherein R' is selected from H, alkyl,
alkenyl, alkynyl, acyl or aryl;

10

$-C(O)R''$, wherein R'' is selected from H, alkyl,
substituted alkyl, alkoxy, alkylamino,
alkenyl, substituted alkenyl, alkynyl,
substituted alkynyl, aryl, substituted
aryl, aryloxy, arylamino, alkylaryl,
substituted alkylaryl, arylalkyl,
substituted arylalkyl, heterocyclic,
substituted heterocyclic or
trifluoromethyl,

15

$-CO_2R'''$, wherein R''' is selected from H,
alkyl, alkenyl or alkynyl;

$-SR'$, $-S(O)R'$, $-S(O)_2R'$ or $-S(O)_2NHR'$, wherein
each R' is as defined above.

7. A method according to claim 5 wherein:

X of Formula I is $-\text{CRR}-\text{C}(\text{R})=\text{C}(\text{R})-(\text{CRR})_m-\text{Z}$,
wherein:

each R is independently selected from
hydrogen, lower alkyl, substituted
lower alkyl, hydroxy, alkoxy (of a
lower alkyl group), halogen,
trifluoromethyl, amino, carboxyl, or
sulfonyl,

m falls in the range of 1 up to 6, and
Z is selected from $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OAc}$, $-\text{CO}_2\text{H}$,
 $-\text{CO}_2\text{Me}$ or $-\text{CO}_2\text{Et}$; and

Y of Formula I is selected from:

$=\text{C}(\text{R})-\text{C}(\text{R})=\text{C}(\text{R})-(\text{CRR})_{n'}-\text{Z}'$ (II),

$=\text{C}(\text{R})-\text{CRR}-\text{CR}(\text{R}')-(\text{CRR})_{n'}-\text{Z}'$ (III), or

$-\text{C}(\text{R})=\text{C}(\text{R})-\text{CR}(\text{R}')-(\text{CRR})_{n'}-\text{Z}'$ (IV), wherein
each R is independently as defined
above,

each R' is independently selected
from H, lower alkyl, substituted
lower alkyl, or a leaving group,
Z' is selected from H, lower alkyl or
substituted lower alkyl, and
n' falls in the range of 1 up to 6.


8. A method according to claim 7 wherein Y of
Formula I is

$=\text{C}(\text{R})-\text{C}(\text{R})=\text{C}(\text{R})-(\text{CRR})_{n'}-\text{Z}'$ (II),

wherein each R is selected from hydrogen, lower alkyl or
substituted lower alkyl, n is 1, n' falls in the range of
about 2 up to 6, and Z' is selected from hydrogen or
lower alkyl.

09155252.092198
86T260"2525T60

9. A method according to claim 7 wherein Y of Formula I is



5 wherein each R is selected from hydrogen, lower alkyl or substituted lower alkyl, R' is selected from hydrogen, lower alkyl, or an hydroxy group, n is 1, n' falls in the range of about 2 up to 6, and Z' is selected from hydrogen or lower alkyl.

10. A method according to claim 5 wherein A is 9-OH, Y is IV, each R is hydrogen, R' is hydroxy, Z is $-CO_2H$, m = 3, Z' is methyl, n = 1 and n' = 4.

11. A method according to claim 5 wherein A is absent, Y is IV, each R is hydrogen, R' is hydroxy, Z is $-CO_2H$, m is 3, Z' is methyl, n = 1 and n' = 4.

12. A method according to claim 5 wherein A is absent, Y is II, each R is hydrogen, R' is hydroxy, Z is $-CO_2H$, m = 3, Z' is methyl, n = 1 and n' = 4.

13. A method according to claim 5 wherein A is absent, Y is I, each R is hydrogen, Z is $-CO_2H$, m = 3, Z' is methyl, n = 1 and n' = 4.

14. A method according to claim 1 wherein said process mediated by PPAR- γ is cell differentiation to produce lipid-accumulating cells.

15. A method according to claim 1 wherein said process mediated by PPAR- γ is the response of the recipient to insulin.

09155252-002498

Sub 35

16. A method of testing a compound for its ability to regulate transcription-activating effects of a peroxisome proliferator activated receptor-gamma (PPAR- γ), said method comprising assaying for changes in the level of reporter protein present as a result of contacting cells containing said receptor and reporter vector with said compound;
- 5
- wherein said reporter vector comprises:
- (a) a promoter that is operable in said cell,
- 10
- (b) a hormone response element, and
- (c) a DNA segment encoding a reporter protein,
- wherein said reporter protein-encoding DNA segment is operatively linked to said promoter for transcription of said DNA segment, and
- 15
- wherein said hormone response element is operatively linked to said promoter for activation thereof.
- 20

17. A method according to Claim 16 wherein said hormone response element is a direct repeat of two or more half sites separated by a spacer of one nucleotide, wherein said spacer can be A, C, G or T,
- 5
- wherein each half site comprises the sequence

-RGBNNM-,

wherein

- R is selected from A or G;
- B is selected from G, C, or T;
- 10
- each N is independently selected from A, T, C, or G; and
- M is selected from A or C;

with the proviso that at least 4 nucleotides of said -RGBNNM- sequence are identical with the nucleotides

15

at corresponding positions of the sequence -AGGTCA-; and

09152222.092498

wherein said response element is optionally preceded by N_x , wherein x falls in the range of 0 up to 5.

Sub 3
18. A method according to claim 17 wherein said response element has at least one copy of the minimal sequence:

AGGACA A AGGTCA,

5 wherein said minimal sequence is optionally flanked by additional residues.

19. A method according to claim 17 wherein said response element has at least one copy of the sequence:

GGACC AGGACA A AGGTCA CGTTC.

Sub 36
20. A method according to claim 16 wherein said compound is a putative antagonist for said peroxisome proliferator activated receptor-gamma, and wherein said contacting is carried out in the presence of

5 increasing concentrations of said compound, and

a fixed concentration of at least one agonist for said peroxisome proliferator activated receptor-gamma.

21. A method according to Claim 16 wherein said contacting is carried out in the further presence of at least one PPAR- γ -selective modulator.

22. A method for preventing obesity, said method comprising administering to a subject in need thereof an amount of a peroxisome proliferator activated receptor-gamma (PPAR- γ) antagonist effective to block
5 cell differentiation to produce lipid-accumulating cells.

091553-092498
051250-252550

23. A method for treating diabetes, said method comprising administering to a subject in need thereof an amount of a peroxisome proliferator activated receptor-gamma (PPAR- γ) agonist effective to lower the blood glucose level of said subject.

add B1

add H1

Add H1

09150000 092198